

Table 1. Survey respondent characteristics (N = 2398)

Characteristic	Percentage (number) of survey respondents
<i>Female gender</i>	87.2% (2091)
<i>Thyroid cancer type</i>	
Any differentiated	87.1% (2088)
Papillary	65.0% (1558)
Follicular	8.7% (209)
Mixed papillary/follicular	11.7% (281)
Hürthle cell	1.7% (40)
Medullary	4.4% (105)
Familial	1.0% (25)
Anaplastic	0.5% (11)
Other	3.4% (81)
Don't know	4.7% (113)
<i>Time since diagnosis</i>	
<1 yr	16.4% (394)
1-5 yrs	55.0% (1318)
6-10 yrs	17.8% (427)
11-15 yrs	5.5% (133)
≥15 yrs	5.3% (126)
<i>Known persistent disease</i>	19.6% (470)
<i>Known distant metastases</i>	14.1% (338)
<i>Familial history of thyroid cancer</i>	12.8% (307)
<i>Age at thyroid cancer diagnosis</i>	
≤19 yrs	4.3% (103)
20-29 yrs	18.2% (436)
30-39 yrs	32.2% (771)
40-49 yrs	24.9% (598)
50-59 yrs	14.7% (353)
60-69 yrs	4.9% (117)
≥70 yrs	0.8% (20)

Percentages may not add up exactly to totals or subtotals due to rounding.