

Table 2. Characteristics of subclinical hypothyroid and healthy control women

	Subclinical hypothyroidism	Healthy control women	p
n	82	51	
Age (yrs)	59.17±7.07	59.69±5.72	0.661
BMI (m/kg ²)	27.89±3.46	27.68±4.66	0.760
Fat mass	42.81±6.76	43.41±5.79	0.649
Menopause	48.05±4.09	48.53±4.58	0.531
Current smoking	16 (19.5%)	10 (19.6%)	0.989
Diabetes mellitus	4 (4,9%)	2 (3.9%)	0.796
Parents fracture	17 (21.0%)	5 (9.8%)	0.093
TSH (mU/liter)	8±2.82	2,23±0,94	<0.001
ft4 (mU/liter)	12.34±2.05	13.82±1.31	<0.001
TPOAb	38 (48.7%)	2 (3.9%)	<0.001
Lumbar spine T score	-1.24±1.10	-1.13 ± 1.59	0.633
Hip T score	-1.08±0.73	-0.68 ±0.90	0.005
Osteocalcin	23.99±12.63	21.79±5.34	0.484
Cross laps	365.76±184.84	306.88±110.73	0.211
Frax-main	6.50±1,58	4.35±1.56	0.001
Frax-hip	1.11±1.94	0.50±0.46	0.030
Previous fractures	18 (22%)	1 (2%)	0.001