Table 1. Disorders in which ghrelin alterations have been found.

 Green upward arrows indicate increased ghrelin levels, red downward arrows indicate decreased ghrelin levels.⁶⁻¹⁸

Condition	Ghrelin levels
Ulcerative colitis ¹	
Crohn's disease ¹	
Poor growth in children ²	
Helicobacter pylori infection ³	V
Takayasu arteritis⁴	V
Coeliac disease⁵	
Cystic fibrosis ⁶	
Coronary artery disease ⁷	V
Hypertension [®]	•
Diabetes (type II) [®]	
Polycystic ovarian syndrome (PCOS) ⁹	V
Cushing's syndrome ¹⁰	V
Hyperthyroidism ¹¹	V
Acromegaly ¹²	V
Gastroesophageal reflux disease (GORD) ¹³	•

- ¹ Ates et al (2008)
- ² Pinsker et al (2011)
- ³ Nweneka & Prentice (2011)
- 4 Yilmaz et al (2012)
- ⁵ Selmiloglu et al (2006)
- ⁶ Monajemzadeh et al (2013)
- 7 Kadoglou et al (2010)

- 8 Poykko et al (2003)
- ⁹ Scholf et al (2002)
- 10 Libe et al (2005)
- ¹¹ Altinova et al (2006)
- 12 Freda et al (2003)
- 13 Nahata et al (2012)