

Table 1. Clinical characteristics of patients with normal pituitary function (NPF), those with isolated pituitary hormone abnormality (IPHA) and those with multiple pituitary hormone abnormalities (MPHA)

	NPF	IPHA	MPHA
N	47	39	7
Sex (male/female)	17/30	11/28	2/5
Age (years)	49.3±1.4	45.5±1.7	54.8±1.7 ^d
BMI (kg/m ²)	23.9±0.6	25.6±0.7 ^b	25.5±2.3 ^b
Time since SAH (yrs)	1.9±0.2	1.7±0.2	1.8±0.4
Vasospasm (yes/no)	2/45	13/26 ^a	3/4 ^a
Hydrocephalus (yes/no)	2/45	5/34	2/5
Epilepsy (yes/no)	4/43	0/39	0/7
GOS score	4.7±0.1	4.4±0.1 ^c	4.6±0.2
Hypertension (yes/no)	24/23	23/16	2/5
Diabetes mellitus (yes/no)	4/43	1/38	0/7
Smoking (yes/no)	29/18	25/14	3/4

^ap <0.001 IPHA and MPHA vs. patients with NPF; ^bp <0.05 IPHA and MPHA vs. patients with NPF; ^cp <0.05 IPHA vs. patients with NPF; ^dp <0.05 MPHA vs. IPHA Descriptive statistics for continuous variables are presented as mean value ± SE.