

Table 1. Pertinent clinical and hormonal data of the subjects studied.

Case	Age yrs	Age at diagnosis yrs	Histology (1)	Stage (2)	BMI kg/m ²	L-T4 µg/week	FT3 pmol/l	FT4 pmol/l	TSH mIU/l	Tg µg/l
Pre-menopausal women										
1	28	18	PTC	1	20.7	700	3.5	16.3	1.52	<0.5
2	31	28	PTC	1	22.1	875	7.1	29.6	<0.03	<0.5
3	33	29	F var PTC	1	19.2	850	4.8	17.4	0.03	<0.5
4	40	38	PTC	1	18.5	1050	5.4	22.5	0.12	<0.5
5	42	41	PTC	1	25.7	1100	5.5	28.6	0.10	<0.5
6	43	18	PTC	1	35.6	1212	4.6	22.4	<0.03	<0.5
7	49	45	PTC	4	35.1	700	4.4	14.3	1.64	<0.5
Post-menopausal women										
8	53	51	PTC	2	27.7	875	3.4	15.1	0.60	<0.5
9	53	44	F var PTC	2	26.8	700	3.1	19.8	0.21	<0.5
10	55	54	F var PTC	2	24.8	787	3.7	23.8	0.92	<0.5
11	57	56	F var PTC	1	22.9	1000	5.2	18.8	1.13	0.5
12	58	56	PTC	1	25.6	700	3.1	13.2	2.31	<0.5
13	58	54	PTC	2	26.9	700	4.1	16.9	0.10	<0.5
14	58	46	PTC	2	25.9	875	4.4	19.0	0.05	<0.5
15	65	62	PTC	1	27.6	1050	4.3	21.9	<0.03	<0.5
16	68	66	PTC	1	36.9	1050	5.2	30.0	<0.03	0.5
17	71	68	Insular	3	32.4	875	4.6	21.4	0.12	<0.5
Males										
18	32	30	FTC	1	22.2	925	6.0	25.2	0.05	0.5
19	38	35	PTC	1	29.4	1500	3.4	14.3	0.62	<0.5
20	50	47	PTC	2	34.1	350	3.1	14.9	3.90	<0.5
21	54	50	PTC	3	23.3	525	5.1	14.4	2.83	<0.5
22	58	56	PTC	3	25.6	700	3.1	13.2	2.31	<0.5
23	61	58	PTC	1	26.2	925	4.1	22.1	9.34	<0.5
24	62	60	PTC	1	25.6	1100	4.3	30.4	2.85	<0.5

¹PTC = papillary thyroid carcinoma, F var PTC = follicular variant of PTC, FTC = follicular thyroid carcinoma.

Thyroid staging was evaluated according to the TNM 2002 classification.